

Quasi-uniform approximation for the radial Schrödinger equation with power-law potentials

V. V. Kudryashov and Yu. V. Vanne

B.I.Stefanov Institute of Physics, National Academy of Sciences of Belarus

F.Skaryna avenue 68, 220072 Minsk, Belarus

E-mail: kudryash@dragon.bas-net.by

Abstract

The uniformly valid approximation to solutions of the radial Schrödinger equation with power-law potentials are obtained by means of the explicit summation of the leading constituent WKB series.

PACS numbers: 02.30.Lt; 03.65.Sq

Key words: radial Schrödinger equation, WKB series, approximate wavefunctions

Similar questions arise when we use the perturbation theory or the variational method in order to solve the radial Schrödinger equation

$$\frac{d^2\Psi(r)}{dr^2} = \frac{Q(r)}{\hbar^2}\Psi(r), \quad Q(r) = 2m \left(V(r) + \frac{\hbar^2 l(l+1)}{2mr^2} - E \right). \quad (1)$$

How to find the unperturbed Hamiltonian or how to find the trial function for an arbitrary given potential? The universal answers are absent. In this sense both mentioned methods are not complete. At the same time the WKB approximation is directly determined by a given potential. The WKB approach deals with a logarithmic derivative

$$Y(r) = \frac{d \ln \Psi(r)}{dr}, \quad \Psi(r) = \exp \left(\int^r Y(r') dr' \right)$$

which satisfies the nonlinear Riccati equation

$$\frac{dY(r)}{dr} + (Y(r))^2 = \frac{Q(r)}{\hbar^2}. \quad (2)$$

In this approach two independent solutions $Y^\pm(r)$ of the Riccati equation are represented by their asymptotic expansions

$$Y_{as}^\pm(r) = \hbar^{-1} \left(\pm Q^{1/2} + \sum_{n=1}^{\infty} \hbar^n Y_n^\pm(r) \right) \quad (3)$$

in powers of Plank's constant $\hbar$. The usual WKB approximation contains a finite number of leading terms $Y_n^\pm(r)$ from the complete expansion $Y_{as}^\pm(r)$. The conventional WKB approximation is not valid at turning points.

As it is well known, the WKB series is divergent. The direct summation of a divergent series does not exist. By summing one means finding a function to which this series is the asymptotic expansion [1]. There are several investigations on properties of the WKB terms [2, 3]. In recent years many studies have been devoted to extracting some useful information about the exact eigenfunctions from the divergent WKB series (see for instance [4] and references therein). In our previous work [5] we reconstructed the WKB series as a sum of new constituent (partial) asymptotic series. The explicit summation of the leading constituent series was performed. As a result the new quasi-uniform approximation to logarithmic derivatives was obtained. This approximation was derived formerly [6] by other means. Our approximation reproduces the known satisfactory approximation [1] near turning points. The proposed method was verified in

the case of the one-dimensional Schrödinger equation without singularities [7]. In the present work we apply our approach to the radial Schrödinger equation with the centrifugal term which is singular at the origin ($r = 0$).

The second-order quasi-uniform approximation has the following form

$$Y_{ap}(r) = Y(r; t) = b_1(r)y_1(a; t) + b_2(r)y_2(a; t) \quad (4)$$

where

$$a(r) = \frac{1}{\hbar^{2/3}} \frac{Q(r)}{|Q'(r)|^{2/3}}, \quad b_1(r) = \frac{1}{\hbar^{2/3}} \frac{Q'}{|Q'|^{2/3}}, \quad b_2(r) = \frac{Q''}{Q'}. \quad (5)$$

Here we introduce notations for logarithmic derivatives

$$y_1(a; t) = \frac{d}{da} \ln (\text{Ai}(a) + t\text{Bi}(a)) \quad (6)$$

of linear combinations of well studied Airy functions $\text{Ai}(a)$ and $\text{Bi}(a)$ [8]. The value of an arbitrary parameter t determines a selected particular solution. We also use notation

$$y_2(a; t) = \frac{1}{30} \left[-8a^2(y_1(a; t))^2 - 4ay_1(a; t) + 8a^3 - 3 \right]. \quad (7)$$

We consider power-law potentials

$$V(r) = \alpha r^k, \quad \alpha > 0, \quad k > 0. \quad (8)$$

When $l = 0$ we deal with the usual one-dimensional problem for odd states which was solved formerly. When $l \neq 0$ we have a new problem. In this case there are two turning points r_- and r_+ ($r_+ > r_-$) which satisfy equation $Q(r) = 0$. Turning points separate three regions.

In the first region where $r > r_+$ we must describe only the decreasing solution of the Schrödinger equation. Therefore in this case we select the unique particular solution $Y(r; 0)$ of the Riccati equation.

In the second region where $r_- < r < r_+$ we must describe oscillating solutions of the Schrödinger equation. Therefore in this case we select two particular solutions $Y(r; \pm i)$ of the Riccati equation.

In the third region where $r < r_-$ we must reproduce the known behaviour of the exact solution $\Psi_{ex}(r)$ of the Schrödinger equation when $r \rightarrow 0$. In this case we have following simple formulas

$$Q(r) \rightarrow \frac{\hbar^2 l(l+1)}{r^2}, \quad \Psi_{ex}(r) \rightarrow r^{l+1}, \quad Y_{ex}(r) \rightarrow \frac{l+1}{r}. \quad (9)$$

At the same time we can derive relations

$$a(r) \rightarrow a_l = \left(\frac{l(l+1)}{4} \right)^{1/3}, \quad b_1(r) \rightarrow -\frac{2a_l}{r}, \quad b_2(r) \rightarrow -\frac{3}{r}. \quad (10)$$

Thus, the approximate logarithmic derivative tends as follow

$$Y(r; t) \rightarrow \frac{c(l, t)}{r} \quad (11)$$

where

$$c(l, t) = -(2a_l y_1(a_l; t) + 3y_2(a_l; t)). \quad (12)$$

Naturally we must demand

$$c(l, t) = l + 1. \quad (13)$$

It is the quadratic algebraic equation for determining the value of t . One of two solutions leads to an unphysical singularity. The second solution (nearest to zero) t_l gives dependence of t on l . We present two numerical examples $t_1 = -0.00393839, t_2 = -0.000523169$. Hence, in third region we select the unique particular solution $Y(r; t_l)$ of the Riccati equation.

Matching particular solutions at turning points we obtain the continuous approximate wavefunction $\Psi_{ap}(r)$ which is represented by following formulas

$$\Psi_1(r) = N \cos \phi_l \exp \left(- \int_r^{r_-} Y(r'; t_l) dr' \right) \quad (14)$$

when $r < r_-$,

$$\begin{aligned} \Psi_2(r) = N \exp \left(\int_{r_-}^r \frac{Y(r'; +i) + Y(r'; -i)}{2} dr' \right) \\ \times \cos \left(\int_{r_-}^r \frac{Q'}{|Q'|} \frac{Y(r'; +i) - Y(r'; -i)}{2i} dr' - \phi_l \right) \end{aligned} \quad (15)$$

when $r_- < r < r_+$ and

$$\Psi_3(r) = \frac{N}{2} (-1)^n \exp \left(\int_{r_-}^{r_+} \frac{Y(r'; +i) + Y(r'; -i)}{2} dr' \right) \exp \left(\int_{r_+}^r Y(r'; 0) dr' \right) \quad (16)$$

when $r > r_+$. Here

$$\phi_l = \frac{\pi}{3} - \arctan t_l \quad (17)$$

and N is a normalization constant.

We have the new quantization condition

$$\int_{r_-}^{r_+} \frac{Q'}{|Q'|} \frac{Y(r; +i) - Y(r; -i)}{2i} dr = \pi \left(n + \frac{1}{3} \right) + \phi_l, \quad n = 0, 1, 2, \dots \quad (18)$$

which determines the spectral value $E_{sp}(n, l)$ of energy implicitly. We denote the approximate wavefunctions with $E = E_{sp}(n, l)$ as $\Psi_{ap}(r; n, l)$.

So the approximate eigenfunctions are defined completely. But the question arises about the optimal approximate eigenvalues because the value $E_{sp}(n, l)$ is not unique choice.

Since explicit expressions for wavefunctions are already obtained we are able to calculate expectation values

$$\bar{E}(n, l) = \langle \Psi_{ap}(n, l) | \hat{H}_r | \Psi_{ap}(n, l) \rangle \quad (19)$$

of the radial Hamiltonian

$$\hat{H}_r = -\frac{\hbar^2}{2m} \frac{d^2}{dr^2} + V(r) + \frac{\hbar^2 l(l+1)}{2mr^2} \quad (20)$$

where $|\Psi_{ap}(n, l)\rangle$ is the vector in the Hilbert space which corresponds to the function $\Psi_{ap}(r; n, l)$.

In accordance with the eigenvalue problem $\hat{H}|\Psi\rangle = E|\Psi\rangle$ we construct the discrepancy vector

$$|D(e; n, l)\rangle = \hat{H}_r |\Psi_{ap}(n, l)\rangle - e |\Psi_{ap}(n, l)\rangle \quad (21)$$

with an arbitrary parameter e while $\hat{H}_r$ and $|\Psi_{ap}(n, l)\rangle$ are given. It is natural to require that the discrepancy vector should not contain a component proportional to the approximate eigenvector. In other words we consider the orthogonality condition $\langle \Psi_{ap}(n, l) | D(e; n, l) \rangle = 0$ as a criterion for selection of the optimal approximate eigenvalue. As a result we get just the expectation value $\bar{E}(n, l)$ while $E_{sp}(n, l)$ does not fulfil the above requirement. It should be also noted that the scalar product $\langle D(e; n, l) | D(e; n, l) \rangle$ is minimized at $e = \bar{E}(n, l)$.

Now we must verify our approximation numerically for the exactly solvable problems. We shall compare the normalized approximate wavefunctions $\Psi_{ap}(r; n, l)$ with the normalized exact wavefunctions $\Psi_{ex}(r; n, l)$.

In order to estimate the closeness of two functions $f_1(r)$ and $f_2(r)$ we consider two corresponding vectors $|f_1\rangle$ and $|f_2\rangle$ in the Hilbert space. Then we construct a deviation vector $|\Delta f\rangle = |f_1\rangle - |f_2\rangle$ and a scalar product

$$\langle \Delta f | \Delta f \rangle = \langle f_1 | f_1 \rangle + \langle f_2 | f_2 \rangle - \langle f_1 | f_2 \rangle - \langle f_2 | f_1 \rangle. \quad (22)$$

Now we can define the relative deviation

$$\delta f = 1 - \frac{\langle f_1 | f_2 \rangle + \langle f_2 | f_1 \rangle}{\langle f_1 | f_1 \rangle + \langle f_2 | f_2 \rangle} \quad (23)$$

as the numerical estimation of the closeness of two functions. Note that $\delta f = 0$ when $f_1(r) = f_2(r)$.

Thus, we get the following estimation

$$\delta \Psi(n, l) = 1 - \langle \Psi_{ex}(n, l) | \Psi_{ap}(n, l) \rangle \quad (24)$$

in the case of normalized real functions $\Psi_{ex}(r; n, l)$ and $\Psi_{ap}(r; n, l)$. The same numerical comparison may be performed

$$\delta \Psi'(n, l) = 1 - \frac{2 \langle \Psi'_{ex}(n, l) | \Psi'_{ap}(n, l) \rangle}{\langle \Psi'_{ex}(n, l) | \Psi'_{ex}(n, l) \rangle + \langle \Psi'_{ap}(n, l) | \Psi'_{ap}(n, l) \rangle} \quad (25)$$

for first derivatives $\Psi'(r) = d\Psi(r)/dr$. Naturally, we can define analogous estimations for higher derivatives.

Now we compare two functions $\hat{H}_r \Psi_{ap}(r; n, l)$ and $\bar{E}(n, l) \Psi_{ap}(r; n, l)$. This comparison may be performed when we do not know exact solutions. As a result we get the relative discrepancy

$$d(n, l) = \frac{\langle \Psi_{ap}(n, l) | \hat{H}_r^2 | \Psi_{ap}(n, l) \rangle - \bar{E}(n, l)^2}{\langle \Psi_{ap}(n, l) | \hat{H}_r^2 | \Psi_{ap}(n, l) \rangle + \bar{E}(n, l)^2} \quad (26)$$

which is directly connected with the initial Schrödinger equation under consideration.

Finally, we characterize our approximation by the usual relative energy error

$$\delta E(n, l) = \frac{\bar{E}(n, l)}{E_{ex}(n, l)} - 1 \quad (27)$$

where $E_{ex}(n, l)$ is the exact energy value.

Firstly, we estimate our approximation in the case of the oscillator potential $V(r) = \alpha_2 r^2$ for which the exact wavefunctions are well known [9]. Table 1 shows that the quasi-uniform approximation gives fairly accurate results for all considered quantities in the oscillator case.

Table 1. $V(r) = \alpha_2 r^2$.

n, l	$\delta \Psi(n, l)$	$\delta \Psi'(n, l)$	$d(n, l)$	$\delta E(n, l)$
0,0	1.595 10^{-5}	7.118 10^{-5}	6.087 10^{-5}	5.635 10^{-5}
1,0	1.504 10^{-6}	3.098 10^{-6}	7.312 10^{-7}	1.884 10^{-6}
2,0	4.271 10^{-7}	6.391 10^{-7}	8.413 10^{-8}	3.348 10^{-7}
0,1	1.132 10^{-3}	6.625 10^{-3}	9.475 10^{-3}	3.083 10^{-3}
1,1	1.337 10^{-3}	2.898 10^{-3}	7.719 10^{-4}	1.011 10^{-3}
2,1	1.506 10^{-3}	2.891 10^{-3}	5.863 10^{-4}	7.271 10^{-4}
0,2	5.052 10^{-4}	4.877 10^{-3}	4.627 10^{-3}	1.492 10^{-3}
1,2	4.173 10^{-4}	8.886 10^{-4}	1.153 10^{-4}	2.487 10^{-4}
2,2	4.635 10^{-4}	8.646 10^{-4}	8.713 10^{-5}	1.813 10^{-4}

The second example is the linear in r potential $V(r) = \alpha_1 r$. When $l = 0$ the quasi-uniform approximation reproduces the exact result. When $l \neq 0$ the exact wavefunctions are unknown but there are results of numerical solution for some energies [10]. We calculate values of d and δE for these states. Table 2 demonstrates validity of our approximation in the linear case.

Table 2. $V(r) = \alpha_1 r$

n, l	$d(n, l)$	$\delta E(n, l)$
0,1	$6.951 \cdot 10^{-3}$	$2.684 \cdot 10^{-3}$
1,1	$5.357 \cdot 10^{-4}$	$8.230 \cdot 10^{-4}$
0,2	$2.565 \cdot 10^{-3}$	$1.160 \cdot 10^{-3}$
1,2	$8.652 \cdot 10^{-5}$	$2.238 \cdot 10^{-4}$

Hence, the performed reconstruction of the WKB series and subsequent explicit summation of the leading constituent (partial) series yield the satisfactory (qualitative and quantitative) description of wavefunctions in the case of the radial Schrödinger equation with power-law potentials.

References

- [1] C. M. Bender and S. A. Orszag, Advanced Mathematical Methods for Scientists and Engineers (McGraw-Hill, New York, 1978).
- [2] C. M. Bender, S. A. Olaussen and P. S. Wang, Phys. Rev. **D16**, 1748 (1977).
- [3] M. Robnik and V. G. Romanovski, J. Phys. **A33**, 5093 (2000).
- [4] E. Delabaere, H. Dilingier and F. Pham, J. Math. Phys. **38**, 6126 (1997).
- [5] V. V. Kudryashov, Doklady of the National Academy of Sciences of Belarus **42** (No. 6), 45 (1998) (in Russian).
- [6] V. V. Kudryashov, Quantum Systems: New Trends and Methods, eds. Y. S. Kim et al (World Scientific, Singapore, 1997) p. 202.
- [7] V. V. Kudryashov and Yu. V. Vanne, Nonlinear Phenomena in Complex Systems, eds. L. Babichev and V. Kuvshinov (Institute of Physics, Minsk, 2000) p. 161.
- [8] M. Abramovitz and I. A. Stegun, Handbook of Mathematical Functions (Dover, New York, 1970).
- [9] S. Flügge, Practical Quantum Mechanics (Springer-Verlag, Berlin), 1999.
- [10] E. Eichten et al, Phys. Rev. **D17**, 3090 (1978).